

Function of a several variables

Q. What is a function?

Function is a rule

Ex: $y = 2x + 3$
dependent \rightarrow independent variable

\rightarrow Role of dependent and independent variable

Dependent variable \rightarrow Range

Independent variable \rightarrow Domain,

Dimension,

Path

$$y = f(x)$$

1. Domain: x -axis

2. Range: y -axis

$$z = f(x, y)$$

1. Two independent variable

2. Two dimension

3. Domain: xy plane

4. Range: z -axis

\Rightarrow Range is always \perp^r to domain.

Partial Differential Equation

1. $y = f(x)$

$$y' = f'(x)$$

Nature of the path of derivative

2. If the function have a single independent variable, function can be expressed in ordinary derivative.

1. $z = f(x, y)$

If the function have more than one independent variable function can be expressed in partial derivative.

2. Notation

(a) $\frac{\partial z}{\partial x} \Big|_{y \rightarrow \text{constant}}$

(c) $\frac{\partial^2 z}{\partial x^2} \Big|_{y \rightarrow \text{constant}}$

(b) $\frac{\partial z}{\partial y} \Big|_{x \rightarrow \text{constant}}$

(d) $\frac{\partial^2 z}{\partial y^2} \Big|_{x \rightarrow \text{constant}}$



$$(c) \frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$$

$$Q \quad z = \sin \pi x \cos y$$

$$\frac{\partial z}{\partial x} = \frac{\partial (\sin \pi x \cos y)}{\partial x} = \cos \pi x \cos y$$

$$\frac{\partial z}{\partial y} = \sin \pi x \frac{\partial (\cos y)}{\partial y} = -\sin \pi x \sin y$$

$$\frac{\partial^2 z}{\partial y^2} = -\sin \pi x \cos y$$

→ Limit : Function of two variables
 ↳ Approximation

$f(x, y) = \frac{x^2 - y^2}{x - y}$; this function is defined at the values of x and y except $x=0$ and $y=0$, $x=y$.

$$\lim_{(x, y) \rightarrow (0, 0)} f(x, y) = \lim_{(x, y) \rightarrow (0, 0)} \frac{x^2 - y^2}{x - y} \quad \left| \begin{array}{l} x > 0, y > 0 \\ x < 0, y < 0 \end{array} \right.$$

$\lim_{x \rightarrow x_0} f(x) = L \rightarrow$ fixed and unique
 $x \rightarrow x_0 \rightarrow$ fixed and unique

$$f(x) = \frac{x^2 - 2^2}{x - 2}$$

$$\lim_{x \rightarrow 2^+} \frac{(x-2)(x+2)}{x-2} = \lim_{x \rightarrow 2^-} \frac{(x-2)(x+2)}{x-2} = 4$$

∴ $\lim_{x \rightarrow 2^+} x+2 = 4 \rightarrow$ Fixed
 ↳ Fixed

A function $f(x, y)$ have limit L , if for every $\epsilon > 0$, \exists a positive number $\delta > 0$, such that

$$|f(x, y) - L| < \epsilon, \text{ whenever } |x - x_0| < \delta \\ |y - y_0| < \delta$$

Symbolically,

$$\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y) = L$$

→ Independent Path: Choose any path function value must be same.

Example 1: Find the limit of function $f(x, y) = \frac{xy}{x^2 + y^2}$ where $x \rightarrow 0$ and $y \rightarrow 0$.

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{xy}{x^2 + y^2}$$

→ First Method

Choose a path $y = mx$

$$\therefore \lim_{(x, mx) \rightarrow (0, 0)} \frac{x(mx)}{x^2 + m^2x^2} = \frac{m}{1+m^2} \quad \text{As, } m \text{ is constant} \\ \Rightarrow \text{Limit does not exist.}$$

Example 2:

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{x - y}{x + y}$$

Choose a path $y = mx$

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{x - mx}{x + mx} = \frac{1 - m}{1 + m}; \quad \text{Here } m \text{ is constant.}$$

As, function depends on constant. Hence, the limit does not exist.

Example 3:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - x\sqrt{y}}{x^2 + y}$$

Choose a path $y = mx^2$

$$\lim_{(x, mx^2) \rightarrow (0,0)} \frac{x^2 - x\sqrt{mx^2}}{x^2 + mx^2} = \lim_{(x, mx^2) \rightarrow (0,0)} \frac{x^2 (1 - \sqrt{m})}{x^2 (1 + m)} ; m \text{ is constant}$$

As; function is dependent on constant 'm'.

\Rightarrow Limit does not exist.

Example 4:

$$\lim_{(x,y) \rightarrow (0,1)} \tan^{-1}\left(\frac{y}{x}\right) = \frac{+\pi}{2} \left[\begin{array}{l} \text{Not fixed} \\ \text{and not} \\ \text{unique} \end{array} \right]$$

Example 5:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3 y}{x^6 + y^2}$$

Choose a path $y = mx$

$$\lim_{(x, mx) \rightarrow (0,0)} \frac{x^3 (mx)}{x^6 + (mx)^2} = \frac{mx^4}{x^2 (x^3 + m)} = \frac{mx^2}{x^3 + m} = 0.$$

Hence, limit exists.

But, if we choose path $y = mx^3$

$$\therefore \lim_{(x, mx^3) \rightarrow (0,0)} \frac{x^3 mx^3}{x^6 + (mx^3)^2}$$

$$\lim_{x \rightarrow (0,0)} \frac{mx^6}{x^6 (1 + m^2)} = \frac{m}{1 + m^2}$$

As, function is dependent on constant

\Rightarrow Limit does not exist.

Example 7

$$\lim_{(x,y) \rightarrow (0,0)} \frac{e^{xy}}{x^2 + y^2}$$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{e^{xy}}{x^2 + y^2} = \lim_{y \rightarrow 0} \lim_{x \rightarrow 0} \frac{e^{xy}}{x^2 + y^2}$$

$$\lim_{y \rightarrow 0} \frac{1}{y^2} = \infty$$

Hence, Limit does not exist.

Continuity of function of two variables;
A function $f(x, y)$ is said to be continuous at a point (x_0, y_0) , if

1. function $f(x, y)$ defined at $x = x_0$ and $y = y_0$.
2. $\lim_{(x,y) \rightarrow (x_0, y_0)} f(x, y)$ exists
3. $\lim_{(x,y) \rightarrow (x_0, y_0)} f(x, y) = f(x_0, y_0)$

If function $f(x, y)$ is defined in domain D . Let $f(x_0, y_0)$ is also defined in D . Then function $f(x, y)$ is continuous at given point (x_0, y_0) if for given $\epsilon > 0$ however small there exist a positive number $\delta > 0$ such that

$$|f(x, y) - f(x_0, y_0)| < \epsilon, \text{ whenever}$$

$$0 < \sqrt{(x-x_0)^2 + (y-y_0)^2} < \delta$$

Example: Test continuity of

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + 2y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

at the point $(0, 0)$

By the definition

1. Function defined at (x_0, y_0)

Here the given function $f(x, y)$ is defined at the point $x_0 = 0, y_0 = 0$. The value of the function at the point $x = 0, y = 0$ is equal to 0.

2. $\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y)$ exist

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{xy}{x^2 + 2y^2}$$

Choose a path $y = mx$

$$\Rightarrow \lim_{(x, mx) \rightarrow (0, 0)} \frac{x^2 m}{x^2 (1 + 2m^2)} = \frac{m}{1 + 2m^2}$$

As, function is dependent on constant 'm'.

Hence, Limit does not exist.

Test the continuity of

$$f(x, y) = \begin{cases} \frac{x-y}{x+y}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

at the point $(0, 0)$.

By the definition of continuity :

1. Function defined at (x_0, y_0)

\therefore Function is defined at a point $x_0 = 0$ and $y_0 = 0$, the value of the function $f(x, y)$ at the point $x_0 = 0, y_0 = 0$ is equal to 0.

2. $\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y)$ exists

$$\therefore \lim_{(x, y) \rightarrow (0, 0)} \frac{x - y}{x + y}$$

\therefore Choose a path $y = mx$

$$\lim_{(x, mx) \rightarrow (0, 0)} \frac{x(1 - m)}{x(1 + m)} = \frac{1 - m}{1 + m}$$

As, function is dependent on constant
 \Rightarrow Hence, Limit does not exist.

3. $\lim_{(x, y) \rightarrow (x_0, y_0)} f(x, y) = f(x_0, y_0)$

As, $\lim_{(x, y) \rightarrow (0, 0)} f(x, y) = f(0, 0)$

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{x - y}{x + y} = 0$$

$$\text{and } \frac{1 - m}{1 + m} \neq 0$$

\therefore Function is discontinuous at point $(0, 0)$.

Example ; Test the continuity of the function

$$f(x) = \begin{cases} \frac{x^2 - x\sqrt{y}}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

By the definition of continuity :

1. Function defined at (x_0, y_0)

∴ $f(x, y)$ is defined at a point $x=0$ and $y=0$, the value of the $f(x, y)$ at the point $x_0=0, y_0=0$ is equal to 0.

$$2. \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - x\sqrt{y}}{x^2 + y^2}$$

Choose a path; $y = mx^2$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - x^2\sqrt{m}}{x^2 + (mx^2)^2} = 1 - \sqrt{m}$$

As, function depends on m
Hence, Limit does not exist.

$$3. \lim_{(x,y) \rightarrow (x_0, y_0)} f(x, y) = f(x_0, y_0)$$

$$\Rightarrow \text{As, } 1 - \sqrt{m} \neq 0$$

⇒ Function is discontinuous at $x_0=0$ and $y_0=0$.

Partial Differentiation

→ Homogenous Function: A function in which each term have a same degree.

$$z = f(x, y) = x^n + x^{n-1}y + x^{n-2}y^2 + \dots + xy^{n-1} + y^n.$$

Degree is n

$$x^n \left[x + \frac{y}{x} + \left[\frac{y}{x} \right]^2 + \dots + \left[\frac{y}{x} \right]^{n-1} + \left[\frac{y}{x} \right]^n \right]$$

$z = x^n f\left(\frac{y}{x}\right)$. Homogenous function whose degree is n .

Ex 1: Find the degree

$$f(x, y) = \frac{x^3 - y^3}{x^2 + y^2}$$

$$f(x, y) = \frac{x^3 [1 - (y/x)^3]}{x^2 [1 + (y/x)^2]}$$

$$z = x f\left(\frac{y}{x}\right)$$

$\Rightarrow n = 1$. Hence, degree = 1.

Ex 2: Find the degree

$$z = f(x, y) = \frac{x^{5/2} - y^{5/2}}{x^{1/2} + y^{1/2}}$$

$$z = \frac{x^{5/2} (1 - (y/x)^{5/2})}{x^{1/2} (1 + (y/x)^{1/2})}$$

$$z = x^2 f\left(\frac{y}{x}\right)$$

$\Rightarrow n = 2$. Hence degree = 2.

Ex 3: Find the degree

$$z = \frac{y \tan^{-1} y}{x}$$

$$= \frac{xy \tan^{-1}\left(\frac{y}{x}\right)}{x} = x f\left(\frac{y}{x}\right) \Rightarrow n = 1.$$

Ex: Show that the function

$$f(x, y) = \begin{cases} (x+y) \sin\left(\frac{1}{x+y}\right), & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

is continuous at the point $(0, 0)$ but not partially differentiable at point $(0, 0)$.

By the definition of continuity

$$|f(x, y) - f(x_0, y_0)| < \epsilon, \text{ whenever } 0 < \sqrt{(x-x_0)^2 + (y-y_0)^2} < \delta$$

As, $f(x, y) = (x+y) \sin\left[\frac{1}{(x+y)}\right]$

and $f(0, 0) = 0$

$$\Rightarrow |f(x, y) - f(0, 0)| = \left| (x+y) \sin\left(\frac{1}{x+y}\right) - 0 \right|$$


$$= |x+y| \left| \sin \frac{1}{x+y} \right|$$

$$\Rightarrow |x+y| \leq |x| + |y| \leq 2\sqrt{x^2 + y^2} < \epsilon$$

Hence, the given function is continuous at the point $(0, 0)$

$$\left| (x+y) \sin \frac{1}{(x+y)} - f(0, 0) \right| < \epsilon \text{ whenever}$$

$$0 < \sqrt{(x-0)^2 + (y-0)^2} < \delta$$



By the definition of the partial differentiation along x -axis \Rightarrow

$$f(x, y) \xrightarrow{\Delta x} f(x + \Delta x, y)$$

$$\frac{\partial z}{\partial x} \Big|_{y=\text{cons}} = \frac{\partial f(x, y)}{\partial x} \Big|_{y=\text{cons}} = \frac{1}{x} (x, y)$$

$$\Delta z = f(x + \Delta x, y) - f(x, y)$$

$$\frac{\Delta z}{\Delta x} = \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

$$\lim_{\Delta x \rightarrow 0} \frac{\Delta z}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

$$\frac{\partial z}{\partial x} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

$$\therefore f(x + \Delta x, y) - f(x, y) = (x + \Delta x + y) \sin \left[\frac{1}{x + \Delta x + y} \right] - (x + y) \sin \left(\frac{1}{x + y} \right)$$

\therefore At $x = 0$ and $y = 0$

$$\Rightarrow f(x + \Delta x, y) - f(0, 0) = \Delta x \sin \frac{1}{\Delta x} - 0$$

$$\therefore f(\Delta x) = \Delta x \sin \frac{1}{\Delta x}$$

$$\lim_{\Delta x \rightarrow 0} f(\Delta x) = \lim_{\Delta x \rightarrow 0} \Delta x \sin \frac{1}{\Delta x}$$

$$\lim_{\Delta x \rightarrow 0} f(\Delta x) = \begin{cases} +1, & \Delta x > 0 \\ -1, & \Delta x < 0 \end{cases}$$

\therefore As, Limit is not continuous unique.

Hence, Limit does not exist.

$\therefore f(x, y)$ is not partially differentiable along x -axis at $(0, 0)$.

Ex; Show that the function,

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + 2y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$$

not continuous at $(0, 0)$ but partially differentiable $(0, 0)$.

By the definition of continuity

Choose a path

$$y = mx$$

$$\therefore \lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + 2y^2} = \lim_{(x, mx) \rightarrow (0,0)} \frac{mx^2}{x^2(1+2m^2)} = \frac{m}{1+2m^2}$$

As, function dependent on m

\Rightarrow hence, limit does not exist.

$$f(x, y) \xrightarrow{\Delta x} f(x + \Delta x, y)$$

$$\therefore \left. \frac{\partial z}{\partial x} \right|_{y \rightarrow \text{constant}} = \frac{\partial f(x, y)}{\partial x} = f_x(x, y)$$

$$\Rightarrow \lim_{\Delta x \rightarrow 0} \frac{\Delta z}{\Delta x} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

$$\therefore \frac{\partial z}{\partial x} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

$$\Rightarrow f(x + \Delta x, y) - f(x, y) = \frac{(x + \Delta x)y}{(x + \Delta x)^2 + 2y^2} - \frac{xy}{x^2 + 2y^2}$$

At point $(0, 0)$

$$\therefore f(0 + \Delta x, 0) - f(0, 0) = \frac{(\Delta x)0}{(\Delta x)^2} - 0$$

$$\Rightarrow \lim_{\Delta x \rightarrow 0} f(\Delta x) = 0$$

\therefore As, Limit is fixed and unique

\Rightarrow Limit exists

Similarly, with y -axis

$\Rightarrow f(x, y)$ is partially differentiable along x -axis as well as y -axis at point $(0, 0)$.

→ Euler Homogenous Theorem

Statement: If $f(x, y)$ is homogenous function in x and y of degree n , then

$$*** \quad x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$$

→ Proof: Consider a homogenous function
$$z = x^n f\left(\frac{y}{x}\right) \quad \text{--- (1)}$$

Partially differentiate equation (1) w.r.t x , where y is a constant.

$$\therefore \frac{\partial z}{\partial x} = \frac{\partial (x^n f(\frac{y}{x}))}{\partial x}$$

$$\frac{\partial z}{\partial x} = x^n \frac{\partial (f(\frac{y}{x}))}{\partial x} + f(\frac{y}{x}) \frac{\partial x^n}{\partial x}$$

$$\Rightarrow \frac{\partial z}{\partial x} = x^n \frac{\partial (f(\frac{y}{x}))}{\partial (\frac{y}{x})} \cdot \frac{\partial (\frac{y}{x})}{\partial x} + n x^{n-1} f(\frac{y}{x})$$

$$\therefore x \frac{\partial z}{\partial x} = -y x^{n-2} f'(\frac{y}{x}) + n x^{n+0} f(\frac{y}{x}) \quad \text{--- (2)}$$

Again, partially differentiating equation (1) w.r.t y , where x is constant

$$\therefore \frac{\partial z}{\partial y} = \frac{x^n \partial f(\frac{y}{x})}{\partial (\frac{y}{x})} \cdot \frac{\partial (\frac{y}{x})}{\partial y} + f(\frac{y}{x}) \frac{\partial x^n}{\partial y}$$

$$y \frac{\partial z}{\partial y} = y x^{n-1} f'(\frac{y}{x}) \quad \text{--- (3)}$$

On adding (2) and (3) and on using (1),
We get:

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$$

Hence proved.

→ Total differentiation

$$\text{If } z = f(x, y)$$

and

$$x = \phi(t)$$

$$y = \psi(t)$$

then

$$z = f(\phi(t), \psi(t))$$

$$\therefore \frac{dz}{dt} = \frac{\partial z}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial z}{\partial y} \cdot \frac{dy}{dt}$$

→ Composite Function (Chain Rule)

$$\text{If } z = f(x, y)$$

where

$$x = \phi(u, v)$$

$$y = \psi(u, v)$$

then

$$z = f(\phi(u, v), \psi(u, v))$$

$\frac{\partial z}{\partial u}$; where v is constant

$$\begin{array}{l} f \rightarrow x \rightarrow u \\ f \rightarrow y \rightarrow u \end{array}$$

$$\frac{\partial z}{\partial u} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial u}$$

For : $\frac{\partial z}{\partial v}$; where u is constant

$$\begin{array}{l} f \rightarrow x \rightarrow v \\ f \rightarrow y \rightarrow v \end{array}$$

$$\frac{\partial z}{\partial v} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial v}$$

$$Q. \quad \text{If } x = e^{2u} + e^{-2v}$$

$$y = e^{-2u} + e^{2v}$$

then prove that

$$\frac{\partial z}{\partial u} - \frac{\partial z}{\partial v} = z \left[x \frac{\partial f}{\partial x} - y \frac{\partial f}{\partial y} \right]$$

By using Chain Rule:

$$\frac{\partial z}{\partial u} = \frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial u}$$

$$\frac{\partial z}{\partial v} = \frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial v}$$

$$\frac{\partial x}{\partial u} = \frac{\partial (e^{2u} + e^{-2v})}{\partial u} = 2e^{2u}$$

$$\frac{\partial x}{\partial v} = \frac{\partial (e^{2u} + e^{-2v})}{\partial v} = -2e^{-2v}$$

$$\frac{\partial y}{\partial u} = \frac{\partial (e^{-2u} + e^{2v})}{\partial u} = -2e^{-2u}$$

$$\frac{\partial y}{\partial v} = 2e^{2v}$$

$$\therefore x \frac{\partial z}{\partial u} = \left\{ \frac{\partial f}{\partial x} \cdot 2e^{2u} + \frac{\partial f}{\partial y} \cdot (-2e^{-2u}) \right\} (e^{2u} + e^{-2v}),$$

$$\frac{\partial z}{\partial v} = \frac{\partial f}{\partial x} \cdot (-2e^{-2v}) + \frac{\partial f}{\partial y} \cdot 2e^{2v}$$

$$\begin{aligned} \therefore \frac{\partial z}{\partial u} - \frac{\partial z}{\partial v} &= (2e^{2u} + 2e^{-2v}) \frac{\partial f}{\partial x} + (-2e^{-2u} - 2e^{2v}) \frac{\partial f}{\partial y} \\ &= 2 \left\{ x \frac{\partial f}{\partial x} - y \frac{\partial f}{\partial y} \right\} \end{aligned}$$

→ Deduction of Euler's formula

If $z = f(x, y)$

and $f(x, y)$ is a homogenous function with degree 'n', then

$$x \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y \frac{\partial^2 z}{\partial y^2} = n(n-1)z$$

Q. If $u = \cos^{-1} \left(\frac{x^2 + y^2}{x + y} \right)$ then find

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = ?$$

$$\therefore u = \cos^{-1} \left(\frac{x^2 + y^2}{x + y} \right)$$

$$\Rightarrow \frac{x^2 (1 + (y/x)^2)}{x (1 + (y/x))} = \cos u$$

$$\Rightarrow \text{Let } z = \cos u = x f(y/x)$$

Comparing with Homogenous equation

$$\Rightarrow n = 1.$$

Hence,

$$x \cdot \frac{\partial z}{\partial x} + y \cdot \frac{\partial z}{\partial y} = nz$$

By using Chain Rule and Euler Homogenous Theorems

$$z \rightarrow u \rightarrow x$$

$$z \rightarrow u \rightarrow y$$

$$\therefore x \frac{\partial z}{\partial u} \frac{\partial u}{\partial x} + y \frac{\partial z}{\partial u} \frac{\partial u}{\partial y} = 1 \cdot z$$

$$x \cdot \frac{\partial (\cos u)}{\partial u} \cdot \frac{\partial u}{\partial x} + y \frac{\partial (\cos u)}{\partial u} \frac{\partial u}{\partial y} = \cos u$$

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = -\cot u \text{ . ans}$$

Q. If $u = \tan^{-1} \left[\frac{x^{3/2} + y^{3/2}}{x^{1/2} + y^{1/2}} \right]$, then find

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = ?$$

$$\therefore u = \tan^{-1} \left[\frac{x^{3/2} + y^{3/2}}{x^{1/2} + y^{1/2}} \right]$$

$$\therefore \tan u = \frac{x^{3/2} + y^{3/2}}{x^{1/2} + y^{1/2}}$$

$$\text{Let } z = \tan u$$

$$\therefore \text{Let } z = \frac{x^{3/2} [1 + (y/x)^{3/2}]}{x^{1/2} [1 + (y/x)^{1/2}]}$$

$$z = x \left[\frac{1 + (y/x)^{3/2}}{1 + (y/x)^{1/2}} \right]$$

$$z = x f\left(\frac{y}{x}\right)$$

Comparing with

$$z = x^n f\left(\frac{y}{x}\right)$$

$$n = 1.$$

Using Euler's Homogenous Equation ;

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$$

$$x \cdot \frac{\partial(\tan u)}{\partial u} \cdot \frac{\partial u}{\partial x} + y \frac{\partial(\tan u)}{\partial u} \cdot \frac{\partial u}{\partial y} = 1 \times \tan u$$

$$\therefore x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = \frac{\tan u}{\sec^2 u} = \sin^2 u \cos u \quad \text{Ans.}$$

Q. If $z = \sin^{-1}\left(\frac{x}{y}\right) + \sin^{-1}\left(\frac{y}{x}\right)$, then find

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = ?$$

$$\text{As ; } z = \sin^{-1}\left(\frac{x}{y}\right) + \sin^{-1}\left(\frac{y}{x}\right)$$

$$z = y^0 \sin^{-1}\left(\frac{x}{y}\right) + x^0 \sin^{-1}\left(\frac{y}{x}\right)$$

Hence ; $n = 0.$

Using Euler's Homogenous Formula;

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = nz$$

Ass; $n = 0$

$$\Rightarrow x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 0. \quad \text{ans}$$

→ Jacobian (J)

Q. What is the application of the Jacobian?

Ans. Transformation between plane.



Q. How will you construct Jacobian?
With the help of a chain rule

$$\frac{\partial f}{\partial u} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial u}$$

$$\frac{\partial f}{\partial v} = \frac{\partial f}{\partial x} \cdot \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \cdot \frac{\partial y}{\partial v}$$

Proof using Cramer's Rule;

$$J = \frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

Q. If $x = u(1+v)$
and $y = v(1+u)$
Then find J .

$$J = \frac{\partial(x, y)}{\partial(u, v)}$$

$$\therefore J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

$$\therefore \frac{\partial x}{\partial u} = 1+v$$

$$\frac{\partial x}{\partial v} = 0+u$$

$$\frac{\partial y}{\partial u} = 0+v$$

$$\frac{\partial y}{\partial v} = 1+u$$

$$\therefore J = \begin{vmatrix} 1+v & u \\ v & 1+u \end{vmatrix}$$

$$= (1+v)(1+u) - uv$$

$$= 1+u+v+uv - uv$$

$$J = 1+u+v \quad \text{ans}$$

Q. If $x = a \cosh \alpha \cos \beta$
 $y = a \sinh \alpha \sin \beta$
 then find J

$$\cosh^2 \alpha - \sinh^2 \alpha = 1$$

$$\therefore J = \frac{\partial(x, y)}{\partial(\alpha, \beta)} = \begin{vmatrix} \frac{\partial x}{\partial \alpha} & \frac{\partial x}{\partial \beta} \\ \frac{\partial y}{\partial \alpha} & \frac{\partial y}{\partial \beta} \end{vmatrix}$$

$$\therefore \frac{\partial x}{\partial \alpha} = \frac{\partial(a \cosh \alpha \cos \beta)}{\partial \alpha} = a \sinh \alpha \cos \beta$$

$$\frac{\partial x}{\partial \beta} = -a \cosh \alpha \sin \beta$$

$$\frac{\partial y}{\partial \alpha} = a \cosh \alpha \sin \beta$$

$$\frac{\partial y}{\partial \beta} = a \sinh \alpha \cos \beta$$

$$J = \begin{vmatrix} a \sinh \alpha \cos \beta & -a \cosh \alpha \sin \beta \\ a \cosh \alpha \sin \beta & a \sinh \alpha \cos \beta \end{vmatrix}$$

$$J = (a \sinh \alpha \cos \beta)^2 + (a \cosh \alpha \sin \beta)^2$$

$$J = a^2 (\sinh^2 \alpha \cos^2 \beta + \cosh^2 \alpha \sin^2 \beta)$$

$$J = a^2 (\sinh^2 \alpha (1 - \sin^2 \beta) + (1 + \sinh^2 \alpha) \sin^2 \beta)$$

$$= a^2 (\sinh^2 \alpha + \sin^2 \beta) \quad \text{ans.}$$

→ Properties of Jacobian

$$JJ' = 1$$

$$\frac{\partial(x, y)}{\partial(u, v)} \times \frac{\partial(u, v)}{\partial(x, y)} = 1$$

$$\therefore J' = 1/J$$

Q. If $x = r \cos \theta$

$$y = r \sin \theta$$

Show that $JJ' = 1$.

$$J = \frac{\partial(x, y)}{\partial(r, \theta)}$$

$$\therefore \frac{\partial x}{\partial r} = \cos \theta$$

$$\frac{\partial y}{\partial r} = \sin \theta$$

$$\frac{\partial x}{\partial \theta} = -r \sin \theta$$

$$\frac{\partial y}{\partial \theta} = r \cos \theta$$

$$\therefore J = \begin{vmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{vmatrix}$$

$$= r \cos^2 \theta + r \sin^2 \theta = r$$

Now

$$\tan \theta = \frac{y}{x}$$

$$\Rightarrow \theta = \tan^{-1} \left(\frac{y}{x} \right)$$

and $r = \sqrt{x^2 + y^2}$

$$\therefore J' = \frac{\partial(r, \theta)}{\partial(x, y)}$$

$$\frac{\partial x}{\partial x} = \frac{\partial \sqrt{x^2 - y^2}}{\partial x} = \frac{x}{\sqrt{x^2 - y^2}}$$

$$\frac{\partial x}{\partial y} = \frac{-y}{\sqrt{x^2 - y^2}}$$

$$\frac{\partial \phi}{\partial x} = \frac{1}{1 + \left(\frac{y}{x}\right)^2} \cdot \frac{-y}{x^2} = \frac{-y}{x^2 + y^2}$$

$$\frac{\partial \phi}{\partial y} = \frac{x^2}{x^2 + y^2} \cdot \frac{1}{x^2} = \frac{x}{x^2 + y^2}$$

Now:

$$J' = \begin{vmatrix} \frac{-x}{\sqrt{x^2 - y^2}} & \frac{+y}{\sqrt{x^2 + y^2}} \\ \frac{-y}{x^2 + y^2} & \frac{x}{x^2 + y^2} \end{vmatrix}$$

$$J' = \frac{1}{x}$$

$$\Rightarrow JJ' = x \left(\frac{1}{x} \right) = 1 \text{ ans}$$

→ Implicit Form of Jacobian

$$1) F_1(u, v, w, x, y, z) = 0$$

$$F_2(u, v, w, x, y, z) = 0$$

$$F_3(u, v, w, x, y, z) = 0$$

$$J = \frac{\partial(u, v, w)}{\partial(x, y, z)} = (-1)^3 \frac{\partial(F_1, F_2, F_3)}{\partial(x, y, z)} \frac{\partial(F_1, F_2, F_3)}{\partial(u, v, w)}$$

$$J = \frac{\partial (u_1, u_2, u_3, \dots, u_n)}{\partial (x_1, x_2, x_3, \dots, x_n)}$$

$$J = \frac{(-1)^n \frac{\partial (F_1, F_2, \dots, F_n)}{\partial (u_1, u_2, \dots, u_n)}}{\frac{\partial (F_1, F_2, \dots, F_n)}{\partial (u_1, u_2, \dots, u_n)}}$$

Q. If

$$x + y + z = u$$

$$y + z = uv$$

$$z = uvw$$

then find J .

$$J = \frac{\partial (u_1, u_2, u_3, \dots, u_n)}{\partial (x_1, x_2, x_3, \dots, x_n)}$$

$$J = \frac{(-1)^n \frac{\partial (F_1, F_2, \dots, F_n)}{\partial (x_1, x_2, \dots, x_n)}}{\frac{\partial (F_1, F_2, \dots, F_n)}{\partial (u_1, u_2, \dots, u_n)}}$$

Q. 1 f

$$\begin{aligned} x + y + z &= u \\ y + z &= uv \\ z &= uvw \end{aligned}$$

then find J_0

$$F_1(x, y, z, u) = x + y + z - u$$

$$F_2(y, z, u, v) = y + z - uv$$

$$F_3(z, u, v, w) = z - uvw$$

$$J = \frac{\partial (x, y, z)}{\partial (u, v, w)}$$

$$\therefore J = \frac{(-1)^3 \frac{\partial (F_1, F_2, F_3)}{\partial (u, v, w)}}{\frac{\partial (F_1, F_2, F_3)}{\partial (x, y, z)}}$$

$$\therefore \frac{\partial (F_1, F_2, F_3)}{\partial (u, v, w)} = \frac{\partial F_1}{\partial u} \quad \frac{\partial F_2}{\partial v} \quad \frac{\partial F_3}{\partial w}$$

$$\frac{\partial F_2}{\partial u} \quad \frac{\partial F_2}{\partial v} \quad \frac{\partial F_2}{\partial w}$$

$$\frac{\partial F_3}{\partial u} \quad \frac{\partial F_3}{\partial v} \quad \frac{\partial F_3}{\partial w}$$



$$\therefore \frac{\partial F_1}{\partial u} = -1$$

$$\frac{\partial F_2}{\partial u} = -v$$

$$\frac{\partial F_3}{\partial u} = -vw$$

$$\frac{\partial F_1}{\partial v} = 0$$

$$\frac{\partial F_2}{\partial v} = -u$$

$$\frac{\partial F_3}{\partial v} = -uw$$

$$\frac{\partial F_1}{\partial w} = 0$$

$$\frac{\partial F_2}{\partial w} = 0$$

$$\frac{\partial F_3}{\partial w} = -uv$$

$$\Rightarrow J = \begin{vmatrix} -1 & 0 & 0 \\ -v & -u & 0 \\ -vw & -uw & -uv \end{vmatrix}$$

$$= -1 (uv^2 + uw^2)$$

$$= -u^2v \text{ ans}$$

and

$$\frac{\partial (F_1, F_2, F_3)}{\partial (x, y, z)} = \begin{vmatrix} \frac{\partial F_1}{\partial x} & \frac{\partial F_2}{\partial y} & \frac{\partial F_3}{\partial z} \\ \frac{\partial F_1}{\partial y} & \frac{\partial F_2}{\partial x} & \frac{\partial F_3}{\partial z} \\ \frac{\partial F_1}{\partial z} & \frac{\partial F_2}{\partial x} & \frac{\partial F_3}{\partial y} \end{vmatrix}$$

$$\begin{vmatrix} \frac{\partial F_1}{\partial x} & \frac{\partial F_2}{\partial y} & \frac{\partial F_3}{\partial z} \\ \frac{\partial F_1}{\partial y} & \frac{\partial F_2}{\partial x} & \frac{\partial F_3}{\partial z} \\ \frac{\partial F_1}{\partial z} & \frac{\partial F_2}{\partial x} & \frac{\partial F_3}{\partial y} \end{vmatrix}$$

$$\begin{vmatrix} \frac{\partial F_1}{\partial x} & \frac{\partial F_2}{\partial y} & \frac{\partial F_3}{\partial z} \\ \frac{\partial F_1}{\partial y} & \frac{\partial F_2}{\partial x} & \frac{\partial F_3}{\partial z} \\ \frac{\partial F_1}{\partial z} & \frac{\partial F_2}{\partial x} & \frac{\partial F_3}{\partial y} \end{vmatrix}$$

$$\frac{\partial F_1}{\partial x} = 1$$

$$\frac{\partial F_2}{\partial x} = 0$$

$$\frac{\partial F_3}{\partial x} = 0$$

$$\frac{\partial F_1}{\partial y} = 0$$

$$\frac{\partial F_2}{\partial y} = 1$$

$$\frac{\partial F_3}{\partial y} = 0$$

$$\frac{\partial F_1}{\partial z} = 0$$

$$\frac{\partial F_2}{\partial z} = 1$$

$$\frac{\partial F_3}{\partial z} = 1$$

$$\therefore \frac{\partial (F_1, F_2, F_3)}{\partial (x, y, z)} = \begin{vmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{vmatrix}$$

$$= +1(1) - 1(0) + 1(0) = 1$$



$$\therefore J = (-1)^3 \frac{(-u^2v)}{1} = u^2v \text{ ans}$$

Q. Show that

$$JJ' = 1 \quad \text{for}$$

$$x = r \sin \theta \cos \phi$$

$$y = r \sin \theta \sin \phi$$

$$z = r \cos \theta$$

$$J = \frac{\partial(x, y, z)}{\partial(r, \theta, \phi)}$$

$$\frac{\partial x}{\partial r} = \sin \theta \cos \phi$$

$$\frac{\partial y}{\partial r} = \sin \theta \sin \phi$$

$$\frac{\partial z}{\partial r} = \cos \theta$$

$$\frac{\partial x}{\partial \theta} = r \cos \theta \cos \phi$$

$$\frac{\partial y}{\partial \theta} = r \cos \theta \sin \phi$$

$$\frac{\partial z}{\partial \theta} = -r \sin \theta$$

$$\frac{\partial x}{\partial \phi} = -r \sin \theta \sin \phi$$

$$\frac{\partial y}{\partial \phi} = r \sin \theta \cos \phi$$

$$\frac{\partial z}{\partial \phi} = 0$$

$$J = \begin{vmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial \phi} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial \phi} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial \phi} \end{vmatrix}$$

$$\begin{vmatrix} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \cos \theta & -r \sin \theta & 0 \end{vmatrix}$$

$$\begin{vmatrix} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \cos \theta & -r \sin \theta & 0 \end{vmatrix}$$

$$J = \begin{vmatrix} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \cos \theta & -r \sin \theta & 0 \end{vmatrix}$$

$$\begin{vmatrix} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \cos \theta & -r \sin \theta & 0 \end{vmatrix}$$

$$\begin{vmatrix} \sin \theta \cos \phi & r \cos \theta \cos \phi & -r \sin \theta \sin \phi \\ \sin \theta \sin \phi & r \cos \theta \sin \phi & r \sin \theta \cos \phi \\ \cos \theta & -r \sin \theta & 0 \end{vmatrix}$$

$$J = \sin\theta \cos\phi (+\kappa^2 \sin^2\theta \cos\phi) - \kappa \cos\theta \cos\phi (-\kappa \sin\theta \cos\theta \cos\phi) + (-\kappa \sin\theta \sin\phi)(-\kappa \sin^2\theta \sin\phi - \kappa \cos^2\theta \sin\phi)$$

$$J = \kappa^2 \sin^3\theta \cos^2\phi + \kappa^2 \sin\theta \cos^2\theta \cos^2\phi + (-\kappa \sin\theta \sin\phi)(-\kappa \sin\phi)$$

$$J = \kappa^2 \sin^3\theta \cos^2\phi + \kappa^2 \sin\theta \cos^2\theta \cos^2\phi + \kappa^2 \sin\theta \sin^2\phi$$

$$J = \kappa^2 \sin\theta \cos^2\phi (\sin^2\theta + \cos^2\theta) + \kappa^2 \sin\theta \sin^2\phi$$

$$J = \kappa^2 \sin\theta$$

$$\therefore J' = \frac{\partial(\kappa, \theta, \phi)}{\partial(x, y, z)}$$

Squaring and adding x, y and z

$$\Rightarrow \kappa^2 = x^2 + y^2 + z^2$$

$$\kappa = \sqrt{x^2 + y^2 + z^2}$$

→ Functionally dependent and functionally independent.

iff

$J = 0 \Rightarrow$ Functionally dependent
 $J \neq 0 \Rightarrow$ Functionally independent

Determinant \rightarrow Polynomial \rightarrow Equation \rightarrow Roots

$$P(x) = 0 \quad (\text{Equation})$$

$$P(x) = ax^2 + bx + c \Rightarrow \text{Polynomial.}$$

$$P(x) = ax^2 + bx + c = 0 \Rightarrow \text{Equation independent.}$$

Ex: Show that the function

$$u = x - y + z$$

$$v = x + y - z$$

$$w = x^2 + y^2 + z^2 - 2yz \quad \text{is functionally independent}$$

$$\frac{\partial(u, v, w)}{\partial(x, y, z)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} & \frac{\partial v}{\partial z} \\ \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{vmatrix}$$

$$\therefore = \begin{vmatrix} 1 & -1 & 1 \\ 1 & 1 & -1 \\ 2x & 2y + 2z & 2z - 2y \end{vmatrix}$$

$$J = 1(2z - 2y + 2y - 2z) + 1(2z - 2y + 2x) + 1(2y + 2z - 2x)$$
$$J = 4z$$

As; $J \neq 0$;

\Rightarrow J is functionally independent.

→ Taylor's Theorem

Function of two variable :

If $f(x, y)$ is continuous and partially derivative at the point $x=a, y=b$. Then

$$f(x+h, y+k) = f(x, y) + (h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y}) f(x, y) + \frac{1}{2!} (h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y})^2 f(x, y) + \frac{1}{3!} (h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y})^3 f(x, y) + \dots + \frac{1}{(n-1)!} (h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y})^{n-1} f(x, y) + R_n$$

→ Remainder term
→ error

→ Taylor's Theorem is used to :

Get the expansion of a real function.

function is undefined

$$f(x+h, y+k) = f(x, y) + (h \frac{\partial}{\partial x} + k \frac{\partial}{\partial y}) f(x, y) + \frac{1}{2!} \left(h^2 \frac{\partial^2}{\partial x^2} + k^2 \frac{\partial^2}{\partial y^2} + 2hk \frac{\partial^2}{\partial x \partial y} \right) f(x, y) +$$

$$\frac{1}{3!} \left(h^3 \frac{\partial^3}{\partial x^3} + k^3 \frac{\partial^3}{\partial y^3} + 3h^2 \frac{\partial^2}{\partial x^2} \frac{\partial}{\partial y} + 3k^2 \frac{\partial^2}{\partial x} \frac{\partial}{\partial y^2} \right) f(x, y) + \dots + R_n$$

Approximation

Q. By using Taylor's Theorem :

Find the expansion of

$f(x, y) = e^x \cos y$ at the point $(0, 0)$.

$$f(x, y) = e^x \cos y$$

$$\frac{\partial f}{\partial x} = e^x \cos y$$

$$\left. \frac{\partial f}{\partial x} \right|_{(0,0)} = 1$$

$$\frac{\partial f}{\partial y} = -e^x \sin y$$

$$\left. \frac{\partial f}{\partial y} \right|_{(0,0)} = 0$$

$$\frac{\partial^2 f}{\partial x^2} = e^x \cos y$$

$$\left. \frac{\partial^2 f}{\partial x^2} \right|_{(0,0)} = 1$$

$$\frac{\partial^2 f}{\partial y^2} = -e^x \cos y$$

$$\left. \frac{\partial^2 f}{\partial y^2} \right|_{(0,0)} = -1$$

$$\frac{\partial^2 f}{\partial y \partial x} = -e^x \sin y$$

$$\left. \frac{\partial^2 f}{\partial y \partial x} \right|_{(0,0)} = 0$$

$$\frac{\partial^2 f}{\partial x^2} = \frac{\partial f}{\partial y} = e^x \cos y (-e^x \sin y)$$

$$\left. \frac{\partial^2 f}{\partial x^2} \cdot \frac{\partial f}{\partial y} \right|_{(0,0)} = 0$$

$$\frac{\partial^2 f}{\partial y^2} \cdot \frac{\partial f}{\partial x} = -e^x \cos y \cdot e^x \cos y$$

$$\left. \frac{\partial^2 f}{\partial y^2} \cdot \frac{\partial f}{\partial x} \right|_{(0,0)} = -1$$

$$\therefore f(x+h, y+k) = f(x, y) + \left[h \frac{\partial f}{\partial x} + k \frac{\partial f}{\partial y} \right] + \frac{1}{2!} \left[h^2 \frac{\partial^2 f}{\partial x^2} + k^2 \frac{\partial^2 f}{\partial y^2} \right.$$

$$\left. + 2hk \frac{\partial^2 f}{\partial x \partial y} \right] + \frac{1}{3!}$$

$$\left[\frac{h^3 \frac{\partial^3 f}{\partial x^3}}{\partial x^3} + \frac{k^3 \frac{\partial^3 f}{\partial y^3}}{\partial y^3} + 3h^2 \frac{\partial^2 f}{\partial x^2} \frac{\partial f}{\partial y} + 3h \frac{\partial f}{\partial x} \cdot k^2 \frac{\partial^2 f}{\partial y^2} \right] + \dots$$

$$\therefore f(x+h, y+k) = 1 + h + \frac{1}{2!} (h^2 - k^2) + \frac{1}{3!} (h^3 - 3hk^2) + \dots$$

To calculate the value of h and k

$$\Rightarrow x - x_0 = h \quad \text{and}$$

$$y - y_0 = k$$

$$\Rightarrow x - 0 = h$$

$$y - 0 = k$$

$$\Rightarrow f(x+h, y+k) = 1 + x + \frac{(x^2 - y^2)}{2!} + \frac{(x^3 - 3xy^2)}{3!} + \dots$$

→ Maxima and minima of function of two variables

1. The point (a, b) is called relative or local minima if $f(a+h, b+k) \geq f(a, b)$, $\forall h$ and k .

The value of the function $f(x, y)$ at point (a, b) is called relative (local) minima value.

2. The point (a, b) is called relative (local) maximum if $f(a+h, b+k) \leq f(a, b)$, for all h and k .



The value of the function $f(x, y)$ at the point (a, b) is called relative (local) maximum value.

3. Stationary point: The point at which maximum/minimum occur is called point of the extreme or stationary point.

4. Saddle Point: The point at which the function is neither maxima or minima is called Saddle point.

5. Extreme Value: The value of the function at the point of extrem or stationary point is called extreme value.

□ $y = f(x)$



Distance : Terminal point - Initial point

: $C - A$

: Decreasing - Increasing = -ve

Local Maxima


Distance : $C - A$

: Increasing - decreasing

: +ve


: Local Minima

$$z = f(x, y)$$



$$f(x+h, y+k) \leq f(x, y)$$

Local Maxima



$$f(x+h, y+k) \geq f(x, y)$$

Local Minima

Saddle Point

Neither maxima nor minima

→ Necessary condition for maxima and minima

1. $\Delta t - s^2 > 0$

2. If $\Delta t - s^2 > 0$ and $r < 0 \Rightarrow$ Maxima

3. If $\Delta t - s^2 > 0$ and $r > 0 \Rightarrow$ Minima

4. If $\Delta t - s^2 = 0 \Rightarrow$ Saddle Point

→ By the partial differentiation notation:

$$p = \partial z / \partial x$$

$$q = \partial z / \partial y$$

$$r = \partial^2 z / \partial x^2$$

$$s = \partial^2 z / \partial y \partial x$$

$$t = \partial^2 z / \partial y^2$$

→ Critical Point :

$$p = \partial z / \partial x = 0$$

$$q = \partial z / \partial y = 0$$

Q. Calculate minima and maxima for

$$f(x, y) = x^3 y^2 (1 - x - y)$$

$$p = \frac{\partial f}{\partial x} = -x^3 y^2 + 3(1 - x - y)x^2 y^2$$

$$q = \frac{\partial f}{\partial y} = -x^3 y^2 + 2(1 - x - y)x^3 y$$

$$r = \frac{\partial^2 f}{\partial x^2} = -3x^2 y^2 - 3x^2 y^2 + 6(1 - x - y)x y^2$$

$$s = \frac{\partial^2 f}{\partial y \partial x} = -3x^2 y^2 + 6(1 - x - y)x^2 y - 2x^3 y$$

$$t = \frac{\partial^2 f}{\partial y^2} = -2x^3 y - 2x^3 y + 2(1 - x - y)x^3$$

\Rightarrow For critical points

$$p = 0 \text{ and } q = 0$$

$$-x^3 y^2 + 3(1 - x - y)x^2 y^2 = 0 \quad \text{and} \quad -x^3 y^2 + 2(1 - x - y)x^3 y = 0$$

on solving we get ;

$$x = 0, \quad y = 0 \quad \text{and}$$

$$x = 1/2, \quad y = 1/3$$

At $(0, 0)$

$$r = 0$$

$$s = 0$$

$$t = 0$$

$$\Rightarrow \quad r t - s^2 = 0 \quad \Rightarrow \text{Saddle point}$$

At $(1/2, 1/3)$

$$r = -8 \left(\frac{1}{4}\right) \left(\frac{1}{9}\right) + 6 \left(1 - \frac{1}{2} - \frac{1}{3}\right) \frac{1}{2} \left(\frac{1}{9}\right) = -\frac{1}{9}$$

$$s = -7 \left(\frac{1}{8}\right) \left(\frac{1}{3}\right) + 2 \left(1 - \frac{1}{2} - \frac{1}{3}\right) \left(\frac{1}{8}\right) = -\frac{1}{8}$$

$$t = -3 \left(\frac{1}{4}\right) \left(\frac{1}{9}\right) + 6 \left(1 - \frac{1}{2} - \frac{1}{3}\right) \frac{1}{4} \times \frac{1}{3} - 2 \times \frac{1}{8} \times \frac{1}{3} = -\frac{1}{12}$$

$$rt - s^2$$

$$= \frac{1}{9} \times \frac{1}{8} - \frac{1}{144} = \frac{1}{144}$$

As; $rt - s^2 > 0$ and $r < 0$

∴ There exists a maxima at $(1/2, 1/3)$ ans.

Q. Find maxima and minima

$$f(x, y) = 2(x^2 - y^2) - xt + yt$$

$$p = 4x - 4x^3$$

$$q = 4y^3 - 4y$$

$$r = 4 - 12x^2$$

$$s = 0$$

$$t = 12y^2 - 4$$

To find stationary points

$$p = 0 \quad \text{and} \quad q = 0$$

$$x = 0 \quad \text{and} \quad x = \pm 1$$

$$y = 0 \quad \text{and} \quad y = \pm 1$$

∴ Stationary points are;

$$(0, 0), \quad (0, \pm 1)$$

$$(\pm 1, 0), \quad (\pm 1, \pm 1)$$

∴ At $(0, 0)$

$$r = 4$$

$$t = -4$$

$$s = 0$$

As, $rt - s^2 < 0$

⇒ $(0, 0)$ acts as saddle point.

At $(0, \pm 1)$

$$r = 4$$

$$\text{As, } rt - s^2 = 64 > 0$$

$$t = 8$$

and $r > 0$

$$s = 0$$

⇒ $(0, \pm 1)$ acts as minima for $f(x, y)$

$$At (\pm 1, 0);$$

$$r = -8$$

$$t = -4$$

$$s = 0$$

$$As \quad rt - s^2 = 32 > 0$$

$$\text{and } r < 0$$

$\Rightarrow (\pm 1, 0)$ acts as maxima for given function.

$$At (\pm 1, \pm 1)$$

$$r = -8$$

$$t = 8$$

$$s = 0$$

$$As, \quad rt - s^2 = -64 < 0$$

$\Rightarrow (\pm 1, \pm 1)$ is saddle point for given function.

Q. Find extreme value of the function :

$$f(x, y) = \sin x + \sin y + \sin(x+y)$$

$$\therefore p = \cos x + \cos(x+y)$$

$$q = \cos y + \cos(x+y)$$

$$r = -\sin x - \sin(x+y)$$

$$s = -\sin(x+y)$$

$$t = -\sin y - \sin(x+y)$$

For stationary point :

$$p = 0 \quad \text{and} \quad q = 0$$

$$\cos x + \cos(x+y) = 0 \quad \text{and} \quad \cos y + \cos(x+y) = 0$$

\Rightarrow on solving we get

$$\cos x = \cos y$$

$$x = y$$

$$\Rightarrow \cos x + \cos 2x = 0$$

$$\cos x + 2\cos^2 x - 1 = 0$$

$$\Rightarrow x = \pi \quad \text{and} \quad x = \pi/3$$

\Rightarrow stationary points :

$$(\pi/3, \pi/3) \quad \text{and} \quad (\pi, \pi)$$

$$\therefore A \text{ is } (\pi/3, \pi/3)$$

$$r = -\sqrt{3}$$

$$s = \sqrt{3}/2$$

$$t = -\sqrt{3}$$

$$\Rightarrow r^2 - s^2 = 3 - \frac{3}{4} = \frac{9}{4} > 0$$

\therefore As $r^2 - s^2 > 0$ and $r < 0$

\Rightarrow Point $(\pi/3, \pi/3)$ acts as maxima for given function

Extreme value of $f(x, y)$ at $(\pi/3, \pi/3)$:

$$f(\pi/3, \pi/3) = 2 \sin \pi/3 + \sin(2\pi/3) = \sqrt{3} + \frac{\sqrt{3}}{2} \text{ ans}$$

$$A \text{ is } (\pi, \pi)$$

$$r = 0$$

$$s = 0$$

$$t = 0$$

$$\Rightarrow r^2 - s^2 = 0 \quad (\text{Lagrange's Method})$$

\therefore

Q. Calculate the extreme value of the function

$$f(x, y) = \sin x \sin y \sin(x+y)$$

$$p = \frac{\partial f}{\partial x} = \cos x \sin y \sin(x+y) + \sin x \sin y \cos(x+y)$$

$$q = \frac{\partial f}{\partial y} = \sin x \cos y \sin(x+y) + \sin x \sin y \cos(x+y)$$

$$r = \frac{\partial^2 f}{\partial x^2} = -\sin x \sin y \sin(x+y) - \sin x \sin y \cos(x+y) + \cos x \cos y \sin(x+y) + \sin x \cos y \cos(x+y) + \cos x \sin y \cos(x+y) - \sin x \sin y \sin(x+y)$$

$$s = \frac{\partial^2 f}{\partial x \partial y} = \cos x \cos y \sin(x+y) + \sin x \cos y \cos(x+y) + \cos x \sin y \cos(x+y) - \sin x \sin y \sin(x+y)$$

$$t = -\sin x \sin y \sin(x+y) + \sin x \cos y \cos(x+y) + \sin x \cos y \cos(x+y) - \sin x \sin y \sin(x+y).$$

For stationary points;

$$p=0 \text{ and } q=0;$$

$$\cos x \sin y \sin(x+y) + \sin x \sin y \cos(x+y) = 0$$

$$\cos x \sin(x+y) + \sin x \cos(x+y) = 0. \quad -1 \Rightarrow \sin(2x+y) = 0$$

$$\text{or } \sin y = 0.$$

$$\sin x \cos y \sin(x+y) + \sin x \sin y \cos(x+y) = 0$$

$$\sin x \sin(x+y) +$$

$$\cos y \sin(x+y) + \sin y \cos(x+y) = 0 \quad - (2) \Rightarrow \sin(x+2y) = 0$$

$$\text{or } \sin x = 0.$$

Q. Calculate the extreme value of

$$f(x, y) = x^3 + y^3 - 3axy$$

$$\therefore p = \frac{\partial f}{\partial x} = 3x^2 - 3ay$$

$$q = 3y^2 - 3ax$$

$$r = 6x$$

$$s = -3a$$

$$t = 6y$$

For critical points:

$$p = 0 \quad \text{and} \quad q = 0$$

$$3x^2 - 3ay = 0 \quad \text{and} \quad 3y^2 - 3ax = 0$$

$$x^2 - ay = 0 \quad \text{and} \quad y^2 - ax = 0$$

$$x^2 = ay \quad \text{or} \quad \frac{x^2}{y} = \frac{ay}{y}$$

$$\frac{ay}{y} = a$$

$$\left(\frac{x}{y}\right)^2 = \left(\frac{y}{x}\right)$$

$$\left(\frac{x}{y}\right)^3 = 1$$

$$y = \frac{x^2}{a}$$

$$\therefore \frac{x^4}{a^2} = ax$$

$$x^3 = a^3$$

$$\therefore a^2 = ay$$

$$x = a$$

$$y = a$$

\therefore Stationary point (a, a)

\Rightarrow At (a, a) ;

$$r = 6a$$

$$s = -3a$$

$$t = 6a$$

$$\Rightarrow rt - s^2 = 36a^2 - 9a^2 = 27a^2 > 0$$

As $rt - s^2 > 0$ and $r > 0$

$\Rightarrow (a, a)$ acts as minima for $f(x, y)$.

$$\text{Extreme value} = a^3 + a^3 - 3a^3 = -a^3 \text{ min}$$



Lagrange's Multipliers Method.

Let $f(x, y, z)$ with variable x, y, z where these variables are connected to $\phi(x, y, z)$ which is represent subjective to condition for the given function $f(x, y, z)$.

For the stationary point of the function $f(x, y, z)$

$$\frac{\partial f}{\partial x} = 0 \quad ; \quad \frac{\partial f}{\partial y} = 0 \quad \text{and} \quad \frac{\partial f}{\partial z} = 0.$$

By using total differentiation for the function $f(x, y, z)$.

$$\frac{\partial f}{\partial x} \cdot dx + \frac{\partial f}{\partial y} \cdot dy + \frac{\partial f}{\partial z} \cdot dz$$

and using total differentiation for the function $\phi(x, y, z)$

$$\frac{\partial \phi}{\partial x} dx + \frac{\partial \phi}{\partial y} dy + \frac{\partial \phi}{\partial z} dz$$

$$F(x, y, z) = \left[\frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz \right] + \lambda \left[\frac{\partial \phi}{\partial x} dx + \frac{\partial \phi}{\partial y} dy + \frac{\partial \phi}{\partial z} dz \right] = 0$$

$\lambda \rightarrow$ Lagrange's multiplier

$$\frac{\partial f}{\partial x} dx + \lambda \frac{\partial \phi}{\partial x} dx = 0$$

$$\frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} = 0 ; \quad \text{Similarly ;}$$

$$\frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} = 0 \quad \text{and}$$

$$\frac{\partial f}{\partial z} + \lambda \frac{\partial \phi}{\partial z} = 0$$

To find the value of x, y, z and λ .
Calculate eq (1), (2) and (3).

Step 1: $F = f(x, y, z) + \lambda \phi(x, y, z)$

Step 2: $\frac{\partial F}{\partial x}, \frac{\partial F}{\partial y}, \frac{\partial F}{\partial z}$

Step 3: $\frac{\partial F}{\partial x} = 0$; $\frac{\partial F}{\partial y} = 0$; $\frac{\partial F}{\partial z} = 0$

Step 4: Calculate the value of x, y, z from 1, 2 and 3.

Step 5: Substituting the value of x, y and z in subjected to condition $\phi(x, y, z)$.

Step 6: We will get value of λ .

Step 7: Substitute the value of λ in step 5.

Step 8: We will get value of x, y and z .

Step 9: Substitute the value of x, y and z in $f(x, y, z)$.

Example: Find the point upon the plane $ax + by + cz = p$ at which the function $x^2 + y^2 + z^2$ has a minimum. Find this minimum function f .

Given:

$$f = x^2 + y^2 + z^2 \quad \text{--- (1)}$$

$$\phi = ax + by + cz = p \quad \text{--- (2)}$$

Suppose:

$$F = f(x, y, z) + \lambda \phi(x, y, z)$$

$$F = (x^2 + y^2 + z^2) + \lambda (ax + by + cz - p)$$

$$\therefore \frac{\partial F}{\partial x} = 2x + \lambda a \quad \text{--- (5)}$$

$$\frac{\partial F}{\partial y} = 2y + \lambda b \quad \text{--- (6)}$$

$$\frac{\partial F}{\partial z} = 2z + \lambda c \quad \text{--- (7)}$$

For stationary point:

$$\therefore \frac{\partial F}{\partial x} = 0 ; \quad \frac{\partial F}{\partial y} = 0 \quad \text{and} \quad \frac{\partial F}{\partial z} = 0$$

$$\Rightarrow x = -\frac{\lambda a}{2}, \quad y = -\frac{\lambda b}{2} \quad \text{and} \quad z = -\frac{\lambda c}{2}$$

Substituting the values of x, y and z in (2)

$$\Rightarrow a\left(-\frac{\lambda a}{2}\right) + b\left(-\frac{\lambda b}{2}\right) + c\left(-\frac{\lambda c}{2}\right) = p$$

$$\therefore \lambda = \frac{-2p}{a^2 + b^2 + c^2}$$

$$\Rightarrow x = \frac{2ap}{(a^2 + b^2 + c^2)}, \quad y = \frac{2bp}{a^2 + b^2 + c^2}, \quad z = \frac{2cp}{a^2 + b^2 + c^2}$$

Putting x, y and z in $F(x, y, z)$

$$f = \frac{p}{(a^2 + b^2 + c^2)} \quad \text{ans}$$

Q. Find the maximum value of $f(x, y, z) = x^p y^q z^r$ subjected to condition $ax + by + cz = p + q + r$

Given:

$$f = x^p y^q z^r$$

$$\phi = ax + by + cz = p + q + r$$

$$F = x^p y^q z^r + \lambda (ax + by + cz - p - q - r)$$

$$\therefore f = x^p y^q z^r$$

$$\therefore \ln f = p \ln x + q \ln y + r \ln z$$

$$\therefore \frac{\partial f}{\partial x} = p \frac{f}{x}$$

$$\frac{\partial f}{\partial y} = q \frac{f}{y}$$

$$\frac{\partial f}{\partial z} = r \frac{f}{z}$$

$$\Rightarrow \frac{\partial F}{\partial x} = p \frac{f}{x} + \lambda a$$

$$\frac{\partial F}{\partial y} = q \frac{f}{y} + \lambda b$$

$$\frac{\partial F}{\partial z} = r \frac{f}{z} + \lambda c$$

\Rightarrow For stationary point

$$\therefore x = -\frac{p f}{\lambda a}$$

$$y = -\frac{q f}{\lambda b}$$

$$z = -\frac{r f}{\lambda c}$$

$$\therefore f = -\frac{p f}{\lambda} - \frac{q f}{\lambda} - \frac{r f}{\lambda} = p + q + r$$

$$f \Rightarrow \lambda = -f$$

Now ; $x = \frac{p}{a}$

$$y = \frac{q}{b}$$

$$z = \frac{r}{c}$$

$$\Rightarrow f(x, y, z) = \left(\frac{p}{a}\right)^p \left(\frac{q}{b}\right)^q \left(\frac{r}{c}\right)^r \text{ ans}$$

Q. If the temperature T at any point $T(x, y, z)$ in the space is

$$T(x, y, z) = Kxyz^2$$

Find the highest temperature on the surface sphere $x^2 + y^2 + z^2 = a^2$.

Given:

$$f(x, y, z) = Kxyz^2 \quad - 1$$

$$\phi(x, y, z) = x^2 + y^2 + z^2 = a^2 \quad - 2$$

$$\therefore F = f(x, y, z) + \lambda \phi(x, y, z)$$

$$F = Kxyz^2 + \lambda(x^2 + y^2 + z^2 - a^2) \quad - 3$$

$$\therefore \frac{\partial F}{\partial x} = Kyz^2 + 2x\lambda \quad - 4$$

$$\frac{\partial F}{\partial y} = Kxz^2 + 2y\lambda \quad - 5$$

$$\frac{\partial F}{\partial z} = 2Kxyz + 2z\lambda \quad - 6$$

For stationary points:

$$Kyz^2 + 2x\lambda = 0 \quad - 7$$

$$Kxz^2 + 2y\lambda = 0 \quad - 8$$

$$2Kxyz + 2z\lambda = 0 \quad - 9$$

$$\therefore Kyz^2 + 2x^2\lambda = 0$$

$$Kxyz + 2y^2\lambda = 0$$

$$2Kxyz + 2z^2\lambda = 0$$

$$f + 2x^2\lambda = 0 \quad - 10$$

$$f + 2y^2\lambda = 0 \quad - 11$$

$$2f + 2z^2\lambda = 0 \quad - 12$$

Adding 10, 11 and 12

$$\Rightarrow 4f + 2\lambda a^2 = 0$$

$$\lambda = \frac{-4f}{2a^2} = \frac{-2f}{a^2}$$

Putting

$$\therefore \lambda = \frac{-2f}{a^2} \text{ in 7, 8 and 9}$$

$$kyz^2 + 2\left(\frac{-2f}{a^2}\right)x = 0$$

$$kyz^2 = \frac{4fx}{a^2}$$

$$k\cancel{y}z^2 = 4 \times \cancel{k} \cancel{y} \cancel{z}^2 \times x$$

$$x = \frac{+a}{2}$$

Similarly:

$$y = \frac{a}{2}$$

$$\text{and } z = \frac{a}{\sqrt{2}}$$

\therefore Putting x, y and z in $f(x, y, z)$

$$f = xyz^2 = \frac{ka^4}{8} \text{ ans.}$$

Q. Find maximum and minimum distance of point $(3, 4, 12)$ from the sphere $x^2 + y^2 + z^2 = 1$.

$$f(x, y, z) = \sqrt{(x-3)^2 + (y-4)^2 + (z-12)^2}$$

$$\text{and } \phi(x, y, z) = x^2 + y^2 + z^2 = 1$$

$$\therefore F = f(x, y, z) + \lambda \phi(x, y, z)$$

$$F = \left(\sqrt{(x-3)^2 + (y-4)^2 + (z-12)^2} + \lambda \cdot (x^2 + y^2 + z^2 - 1) \right)$$

$$\frac{\partial F}{\partial x} = \frac{1 \times \lambda(x-3)}{\lambda \sqrt{(x-3)^2 + \dots}} + 2\lambda x$$

and so on.

$$\Rightarrow \frac{\partial F}{\partial x} = 0$$

$$\therefore \frac{1(x-3)}{\lambda} + 2\lambda x = 0$$

$$x = \frac{3}{2\lambda f + 1} \quad \text{and}$$

$$y = \frac{4}{2\lambda f + 1} \quad \text{and}$$

$$z = \frac{12}{2\lambda f + 1}$$

and

$$\lambda = 6/f \quad \text{and} \quad \lambda = -7/f$$

↑ Maximum
distance

↑ Minimum
distance